


Legionella

What is Legionella?

Legionella are characterized as gram-negative, aerobic, unencapsulated bacilli. Fifty bacterial species with more than 70 serogroups make up this genus; 20 of the species are known to be associated with human infection. Most notably, this bacterium was identified as the cause of the July 1976 outbreak of pneumonia among people at a convention of the American Legion in Philadelphia. Accordingly, the previously unknown bacteria and the disease it caused, Legionnaires' disease, were named.



What is Legionnaires' disease?

Legionnaires' disease, or Legionellosis, is a severe infection caused by *Legionella* species, primarily *L. pneumophila*. In fact, *L. pneumophila* is responsible for 90% of infections. The disease typically presents as pneumonia and symptoms may include a high fever, chills, cough, muscle aches, headaches, and diarrhea.

Pontiac fever is an acute, febrile, self-limited illness that has been linked to *Legionella* species. This infection is generally characterized by fever, fatigue, and muscle pain. Antimicrobial treatment does not need to be administered to treat this infection, but should be prescribed for cases of Legionnaires' disease.

What are the most common species that cause human infection?

Most reported cases of Legionnaires' disease are caused by *Legionella pneumophila*, serogroup 1. The estimated proportion of cases caused by species is:

- *L. pneumophila* (91.4%)
- *L. micdadei* (2.8%)
- *L. longbeachae* (2.2%)
- *L. dumoffii* (1.5%)
- *L. bozemanii* (1.3%)

Where is Legionella found?

The bacterium is found in the following:

- Environmental water sources
- Potable water, such as that from faucets, showers, and hot water tanks
- Cooling towers and hot tubs

Legionella

Who is at risk of *Legionella* infection?

Legionella infection can occur in a variety of hosts. The incidence of Legionnaires' disease depends on the degree of water contamination, the intensity of the patient's exposure to that water, and the susceptibility of the patient.

The presence of the bacteria in the aforementioned sources does not in and of itself constitute a dangerous condition or disease threat to healthy individuals. *Legionella* in a hospital water system increases the risk of hospital acquired infection, particularly for the immuno-compromised and those with underlying diseases. Transplant patients are at risk for infection. Similarly, cigarette smoking, chronic lung disease, and immunosuppressive treatment increase the risk for disease.

How is *Legionella* transmitted?

Modes of transmission of *Legionella* include the following:

- Aspiration of water contaminated with the organism
- Contaminated respiratory-therapy equipment
- Inhalation of aerosols containing *Legionella*

What are the clinical manifestations of infection?

Symptoms of infection may develop after a few days or up to 2 weeks after exposure to the bacterium. Pneumonia is the major clinical presentation of *Legionella* infection, typically caused by aspiration or aerosolization of water from a contaminated water source. The persistence of pulmonary infiltrates is suggestive of Legionnaires' disease. Other symptoms include fever, headache, confusion, and muscle pain. Some advanced disease symptoms may include gastrointestinal and nervous system problems.

How is *Legionella* infection diagnosed?

Diagnosis of Legionnaire's disease is made based upon the following criteria:

- Culture of the bacteria on selective media
- Detection of *Legionella* antigen in urine (urinary antigen test)
- Detection of *Legionella*-specific antibodies in serum (serology)

It is necessary to conduct antibody testing of both acute- and convalescent-phase sera when diagnosing Legionnaires' disease.

Is infection by the bacterium difficult to treat? What treatment is available?

Antimicrobial agents such as macrolides and quinolones are effective in treating *Legionella* infections. The following are some of the antimicrobials that could be given for oral therapy:

- | | |
|------------------|-----------------|
| • Azithromycin | • Ciprofloxacin |
| • Clarithromycin | • Moxifloxacin |
| • Roxithromycin | • Gemifloxacin |
| • Levofloxacin | • Tetracycline |

There is currently no vaccine available for the disease.

Legionella

How can *Legionella* be monitored and controlled in water supplies?

There are several short term and long term water treatment options to reduce *Legionella* bacteria in water systems. These include chemical treatment or repeated thermal eradication. If illness is suspected to be from *Legionella* in a contaminated water source, the water should be immediately tested. Additionally, experts should be consulted for guidance regarding treatment of the water system.

Sources:

Stout JE, Muder RR, Mietzner S, Wagener MM, et al. role of environmental surveillance in determining risk for hospital-acquired Legionellosis: a national surveillance study with clinical correlations. *Infect Control Hosp. Epidemiol.* 2007;28:818-824.

Yu VL, Stout JE. Community-acquired Legionnaires' disease: Implications for underdiagnosis and laboratory testing. *Clin Infect Dis.* 2008; 46:1365-1367.

www.legionella.org

Disclaimer: The information contained in this "Fact Sheet" is provided for informational purposes only and not for the purpose of providing legal or medical advice. You should contact your physician or attorney to obtain advice with respect to any particular issue or problem.